Corrigendum to: Fatty Degeneration of Subscapularis Muscle after Total Shoulder Arthroplasty: A CT-scan Analysis from SFI study Group

Hugo Barret [1] **(*), Arnaud Godeneche** [2], **Thierry Joudet** [3], **Christophe Charousset** [4], **Stephane Audebert** [5], **Yves Lefebvre** [6], **Johannes Barth** [7], **Jerome Garret** [8], **David Gallinet** [9], **Jacques Guery** [10], **SFI** [11], **Nicolas Bonnevialle** [1]

(*) corresponding author: hugobarret89@gmail.com
1. Hôpital Pierre Paul Riquet - CHU de Toulouse - Place Baylac, 31059, Toulouse Cedex 09, France
2. Centre Orthopédique Santy, Ramsay Santé, Hôpital Privé Jean Memoz, Lyon, France
3. Clinique du Libournais, Libourne, France
4. Institut Ostéo Articulaire Paris Courcelles, Paris, France
5. Department of Orthopaedic Surgery, Clinique du Cambresis, Cambrai, France
6. Institut de l'épaule de Strasbourg (Strasbourg Shoulder Institute), Strasbourg, France
7. Clinique des Cèdres, Echirolles, France
8. Clinique du Parc, Groupe ELSAN, Lyon, France
9. Centre épaule main Besançon, 16, rue Madeleine-Brès, 25000 Besançon, France
10. Polyclinique du Val de Loire, Nevers, France
11. Shoulder Friends Institute, 60 Rue de Courcelles, 75008 Paris, France

A Corrigendum on: Fatty Degeneration of Subscapularis Muscle after Total Shoulder Arthroplasty: A CT-scan Analysis from SFI study Group

by Barret, H., Godeneche, A., Joudet, T., et al. (2025). Fatty Degeneration of Subscapularis Muscle after Total Shoulder Arthroplasty: A CT-scan Analysis from SFI study Group. The Hive Musculoskeletal Journal, 1(2).

https://doi.org/10.70885/hmsj.2026.01.003

The authors regret that Tables 1 and 2 were omitted from the originally published version of this article due to an author oversight. The missing tables have been incorporated into the online versions of the article.

This correction does not affect the results or conclusions of the paper.

The authors apologize for any inconvenience caused.

The online version of this article can be found under DOI: https://doi.org/10.70885/hmsj.2025.08.001

Published online: 8/2025; Volume 1 - 2025, Issue 2

Table 1: Tenotomy versus Peel Off

	SSc Tenotomy N = 77	Peel off N = 27	P value
Upper half fatty infiltration of SSc	2.06 (± 0.8)	2.0 (± 0.8)	0.839
Lower half fatty infiltration of SSc	1.18 (± 0.9)	0.78 (± 0.6)	0.062
SSc Hypertrophic			0.886
Yes	25 (32%)	7 (26%)	
No	52 (68%)	20 (74%)	
Upper half fatty infiltration of SSc			0.311
1	13 (16.9%)	6 (22.2%)	
2	43 (55.8%)	12 (44.5%)	
3	16 (20.8%)	9 (33.3%)	
4	5 (6.5%)	0	
Lower half fatty infiltration of SSc			0.706
0	41 (53.2%)	17 (63%)	
1	25 (32.5%)	7 (25.9%)	
2	4 (5.2%)	2 (7.4%)	
3	7 (9.1%)	1 (3.7%)	

SSc = Subscapularis

Table 2: Transosseous repair vs tendon-tendon repair

	Transosseous repair N = 89	Tendon-tendon repair + anchors N = 15	P value
Mean Upper SSc fatty infiltration	1.4 (± 0.7)	2.2(± 1)	0.04*
Lower SSc Hypertrophic			0.33
Yes	29 (33%)	3 (20%)	
No	60 (67%)	12 (80%)	
Upper SSc infiltration			0.03*
0	36 (41%)	2 (13.0%)	
1	33 (37%)	4 (27.0%)	
2	10 (11%)	4(27%)	
3	8 (9%)	4 (27%)	
4	2 (2%)	1 (6.0%)	

SSc = Subscapularis; * statistically significant result